

ZIMPRO CAV FATTENING



INDICATIONS FO USE:

ZIMPROCAV FATTENING main raw material is **NARASIN**, an exclusive lonophore for use in animal feed, which aims to increase feed efficiency and weight gain in beef cattle. **ZIMPROCAV FATTENING** brings together two performance enhancing additives, **NARASIN** and **VIRGINIAMYCIN**, in addition to a blend of vitamins and Probiotic additives.

HOW TO USE:

- Mix 250 grams of **ZIMPROCAV FATTENING** in 25 kg of mineral supplement and give it freely to the animals, preferably in covered troughs.

- Mix 1.0 kg of **ZIMPROCAV ENGORDA** in one ton (1,000 kg) of feed and give the final product after mixing it. Always maintain the good availability of roughage for animals, whether in the form of pasture and/or alternative roughage such as: fresh chopped sugar cane, fresh chopped napier, silages, grass hay.

BENEFITS:

- Promotes the growth of beneficial bacteria
- Greater energy production
- Higher feed conversion
- Greater productivity and performance each in natura, chopped napier in natura, silages, grass hay



PRESENTATION:

5, 10, 20 and 25 kg buckets

Guarantee levels per kg of product:

Description	Guarantee	Unit
Vitamin A (minimum)	200,000.0000	IU/kg
Vitamin D3 (minimum)	10,000.0000	IU/kg
Vitamin E (minimum)	100.0000	IU/kg
Vanilla Aroma (minimum)	2,000.0000	mg/kg
Crude Protein (minimum)	15.5000	g/kg
Narasin (minimum)	1,200.0000	mg/kg
Virginiamycin (minimum)	1,000.0000	mg/kg
Saccharomyces cerevisiae (minimum)	1.5000x10E12	cfu/kg
Ruminobacter amylophilum (minimum)	6.0000x10E10	cfu/kg
Ruminobacter succinogenes (minimum)	6.0000x10E10	cfu/kg
Succinobacterium dextrinosolvens (minimum)	8.8000x10E10	cfu/kg
Bacillus cereus (minimum)	7.0000x10E10	cfu/kg
Lactobacillus acidophilus (minimum)	7.0000x10E10	cfu/kg
Enterococcus faecium (minimum)	7.0000x10E10	cfu/kg

CONTRAINDICATIONS:

- Do not allow turkeys, horses or other equines to have access to formulations containing Narasin. Ingestion of this product by these species can be fatal.

DRUG INTERACTIONS:

- This product contains lonophore, and its simultaneous use with certain substances may be contraindicated.

GRACE PERIOD:

- **SLAUGHTER:** Slaughter of treated animals should only be done 3 days after the last administration.
- **MILK:** this product should not be applied to females producing milk for human consumption.